

Comparison of Measurements Between Different Projects and Funders

Version 19

F

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Cervical Cancer Screening	<ul style="list-style-type: none"> (Reference: CMS124v9) Between 23 and 64 years of age at the beginning of the measurement period Had at least one medical encounter during measurement period Female (or, patients of all genders who have a cervix) Exclusion: had a hysterectomy and no residual cervix, acquired absence of the cervix, a congenital absence of the cervix, or was receiving hospice care during the measurement period 	<p>Patients who had either of the following:</p> <ul style="list-style-type: none"> Cervical cytology performed in the three years prior to the end of the measurement period (age at least 21 years at the time of the test) Cervical human papillomavirus (HPV) testing performed in the five years prior to the end of the measurement period (age at least 30 years at the time of the test) 	<ul style="list-style-type: none"> Continuously enrolled PHP members between 24 and 64 years of age at the end of the measurement period Female (or, patients of all genders who have a cervix) Exclusion: had a hysterectomy and no residual cervix, acquired absence of the cervix, a congenital absence of the cervix, or was in hospice or palliative care during the measurement period 	<p>Patients who had either of the following:</p> <ul style="list-style-type: none"> Cervical cytology performed in the three years prior to the end of the measurement period (age at least 21 years at the time of the test) Cervical human papillomavirus (HPV) testing performed in the five years prior to the end of the measurement period (age at least 30 years at the time of the test) 				
Breast Cancer Screening	<ul style="list-style-type: none"> (Reference: CMS125v9) Between 51 and 73 years of age at the beginning of the measurement period Had at least one medical encounter during measurement period Female (according to sex at birth) Exclusions: had complete bilateral mastectomy or a right and a left complete unilateral mastectomy; in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients who had a mammogram during the 27 months prior to the end of the measurement period</p>	<ul style="list-style-type: none"> Continuously enrolled PHP members between 52 and 74 years of age at the end of the measurement period Female (according to sex at birth) Exclusions: had complete bilateral mastectomy or a right and a left complete unilateral mastectomy; in hospice or palliative care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients who had a mammogram during the 27 months prior to the end of the measurement period</p>				
Colorectal Cancer Screening	<ul style="list-style-type: none"> (Reference: CMS130v9) Between 50 and 74 years at the beginning of the measurement period Had at least one medical encounter during the measurement period Exclusions: had colorectal cancer; had a total colectomy; in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one of the following in the time-frame indicated:</p> <ul style="list-style-type: none"> Fecal occult blood test (FOBT) in the year prior to the end of the measurement period FIT-DNA in the 3 years prior to the end of the measurement period Flexible Sigmoidoscopy in the 5 years prior to the end of the measurement period Computerized tomography (CT) colonography in the 5 years prior to the end of the measurement period Colonoscopy in the 10 years prior to the end of the measurement period 	<ul style="list-style-type: none"> Continuously enrolled PHP members between 51 and 75 years at the end of the measurement period Exclusions: had colorectal cancer; had a total colectomy; in hospice or palliative care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one of the following in the time-frame indicated:</p> <ul style="list-style-type: none"> Fecal occult blood test (FOBT) in the year prior to the end of the measurement period FIT-DNA in the 3 years prior to the end of the measurement period Flexible Sigmoidoscopy in the 5 years prior to the end of the measurement period Computerized tomography (CT) colonography in the 5 years prior to the end of the measurement period Colonoscopy in the 10 years prior to the end of the measurement period 	<ul style="list-style-type: none"> Between 50 and 74 years at the beginning of the measurement period Had at least one medical encounter during the measurement period Exclusions: had colorectal cancer; had a total colectomy; in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one of the following in the time-frame indicated:</p> <ul style="list-style-type: none"> Fecal occult blood test (FOBT) in the year prior to the end of the measurement period FIT-DNA in the 3 years prior to the end of the measurement period Flexible Sigmoidoscopy in the 5 years prior to the end of the measurement period Computerized tomography (CT) colonography in the 5 years prior to the end of the measurement period Colonoscopy in the 10 years prior to the end of the measurement period 		

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Blood Sugar Control Among Patients With Diabetes	<ul style="list-style-type: none"> (Reference: CMS122v9) Between 18 and 74 years of age at the beginning of the measurement period Diagnosed with diabetes Had at least one medical encounter during measurement period Exclusion: in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<ul style="list-style-type: none"> Patients with an HbA1c lab performed within the year prior to the end of the measurement period and the latest result had a value greater than 9% Note that this measure focuses on the negative outcome 	<ul style="list-style-type: none"> Continuously enrolled PHP members between 18 and 75 years of age at the end of the measurement period Diagnosed with diabetes Had at least two outpatient, telephone or virtual encounters on different dates of service during the two years prior to the end of the measurement period Exclusion: in hospice or palliative care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<ul style="list-style-type: none"> Patients with an HbA1c lab performed within the year prior to the end of the measurement period and the latest result had a value equal to or less than 9%. 	<ul style="list-style-type: none"> Between 18 and 74 years of age at the beginning of the measurement period Diagnosed with diabetes Had at least one medical visit during measurement period Exclusion: in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<ul style="list-style-type: none"> Patients with an HbA1c lab performed within the year prior to the end of the measurement period and the latest result had a value equal to or less than 9%. 	<ul style="list-style-type: none"> Between 18 and 74 years of age at the beginning of the measurement period Diagnosed with diabetes Had at least one medical encounter during measurement period Exclusion: in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<ul style="list-style-type: none"> Patients with an HbA1c lab performed within the year prior to the end of the measurement period and the latest result had a value greater than 9% Note that this measure focuses on the negative outcome
Blood Pressure Control Among Patients With Diabetes							<ul style="list-style-type: none"> Between 18 and 74 years of age at the beginning of the measurement period Diagnosed with diabetes Had at least one medical encounter during the measurement period Exclusion: in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<ul style="list-style-type: none"> Patients with at least one blood pressure performed in the measurement period and the latest result had a value less than 140/90 mmHg

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Controlling High Blood Pressure	<ul style="list-style-type: none"> (Reference: CMS165v9) Between 18 and 84 years at the beginning of the measurement period Diagnosed with essential hypertension Had at least one medical encounter during measurement period Exclusions: pregnant during measurement period; with end stage renal disease (ESRD), dialysis, or renal transplant before the end of the measurement period; in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one blood pressure performed within the year prior to the end of the measurement period and the latest result had a value less than 140/90 mmHg</p> <p>Blood pressure readings for this measure can be obtained in the following situations:</p> <ul style="list-style-type: none"> Blood pressure taken by an appropriately trained staff member in a clinical setting Blood pressure readings from a remote device that are digitally stored and transmitted directly to the electronic health record Blood pressure readings from a remote digital device that an appropriately trained staff member can confirm visually during an E-visit or virtual check 	<ul style="list-style-type: none"> Continuously enrolled PHP members between 18 and 85 years at the end of the measurement period Was diagnosed with essential hypertension any time before 6 months prior to the end of the measurement period Had at least two outpatient, telephone or virtual encounters on different dates of service during the two years prior to the end of the measurement period Exclusions: pregnant during measurement period; with end state renal disease (ESRD), kidney transplant or dialysis before the end of the reporting period; in hospice or palliative care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one blood pressure performed within the year prior to the end of the measurement period and the latest result had a value less than 140/90 mmHg</p> <p>Blood pressure readings for this measure can be obtained in the following situations:</p> <ul style="list-style-type: none"> Blood pressure taken by an appropriately trained staff member in a clinical setting Blood pressure readings from a remote device that are digitally stored and transmitted directly to the electronic health record Blood pressure readings from a remote digital device that an appropriately trained staff member can confirm visually during an E-visit or virtual check Self-reported blood pressure readings from a digital device collected verbally from the patient by an appropriately trained staff member during an outpatient visit, telephone visit, e-visit, virtual check-in, or remote monitoring event Self-reported blood pressure readings from a digital device sent to the health center via the patient portal or e-mail 	<ul style="list-style-type: none"> Between 18 and 84 years at the beginning of the measurement period Diagnosed with essential hypertension Had at least one medical visit during measurement period Exclusions: pregnant during measurement period; with end stage renal disease (ESRD), dialysis, or renal transplant before the end of the measurement period; in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one blood pressure performed within the year prior to the end of the measurement period and the latest result had a value less than 140/90 mmHg</p> <p>Blood pressure readings for this measure can be obtained in the following situations:</p> <ul style="list-style-type: none"> Blood pressure taken by an appropriately trained staff member in a clinical setting Blood pressure readings from a remote device that are digitally stored and transmitted directly to the electronic health record Blood pressure readings from a remote digital device that an appropriately trained staff member can confirm visually during an E-visit or virtual check Self-reported blood pressure readings from a digital device collected verbally from the patient by an appropriately trained staff member during an outpatient visit, telephone visit, e-visit, virtual check-in, or remote monitoring event Self-reported blood pressure readings from a digital device sent to the health center via the patient portal or e-mail 	<ul style="list-style-type: none"> Between 18 and 84 years at the beginning of the measurement period Diagnosed with essential hypertension Had at least one medical visit during measurement period Exclusions: pregnant during measurement period; with end stage renal disease (ESRD), dialysis, or renal transplant before the end of the measurement period; in hospice care during the measurement period; aged 66 or older and living long-term in an institution for more than 90 days during the measurement period; aged 66 and older with advanced illness and frailty 	<p>Patients with at least one blood pressure performed within the year prior to the end of the measurement period and the latest result had a value less than 140/90 mmHg</p> <p>Blood pressure readings for this measure can be obtained in the following situations:</p> <ul style="list-style-type: none"> Blood pressure taken by an appropriately trained staff member in a clinical setting Blood pressure readings from a remote device that are digitally stored and transmitted directly to the electronic health record Blood pressure readings from a remote digital device that an appropriately trained staff member can confirm visually during an E-visit or virtual check Self-reported blood pressure readings from a digital device collected verbally from the patient by an appropriately trained staff member during an outpatient visit, telephone visit, e-visit, virtual check-in, or remote monitoring event Self-reported blood pressure readings from a digital device sent to the health center via the patient portal or e-mail
Hypertension Prevalence								

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Statin Therapy for the Prevention and Treatment of Cardiovascular Disease	<ul style="list-style-type: none"> (Reference: CMS347v4) Had at least one medical encounter during the measurement period Three groups of patients quality for being at high risk for cardiovascular events (age calculated at the beginning of the measurement period): <ol style="list-style-type: none"> Patients aged 21 years or older with an active clinical diagnosis of atherosclerotic cardiovascular disease (ASCVD); Patients aged 21 years or older who have ever had an LDL-C result greater than or equal to 190 mg/dL or with a previous or active diagnosis of familial or pure hypercholesterolemia; Patients aged 40 to 75 years with an active clinical diagnosis of diabetes and with any LDL-C result of 70 to 189 mg/dL in the three years prior to the end of the measurement period Exclusions: Patients who were pregnant or breastfeeding in the measurement period; with end-stage renal disease (ESRD) or rhabdomyolysis; with hepatitis A or B, hepatic disease or insufficiency; with allergy or intolerance to statin medication; received palliative 	Patients with documentation demonstrating that statin therapy was being used during the year prior to the end of the measurement period					<ul style="list-style-type: none"> Had at least one medical encounter during the measurement period Three groups of patients quality for being at high risk for cardiovascular events (age calculated at the beginning of the measurement period): <ol style="list-style-type: none"> Patients aged 21 years or older with an active clinical diagnosis of atherosclerotic cardiovascular disease (ASCVD); Patients aged 21 years or older who have ever had an LDL-C result greater than or equal to 190 mg/dL or with a previous or active diagnosis of familial or pure hypercholesterolemia; Patients aged 40 to 75 years with an active clinical diagnosis of diabetes and with any LDL-C result of 70 to 189 mg/dL in the three years prior to the end of the measurement period Exclusions: Patients who were pregnant or breastfeeding in the measurement period; with end-stage renal disease (ESRD) or rhabdomyolysis; with hepatitis A or B, hepatic disease or insufficiency; with allergy or intolerance to statin medication; received palliative care during measurement 	Patients with documentation demonstrating that statin therapy was being used during the year prior to the end of the measurement period
Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antiplatelet	<ul style="list-style-type: none"> (Reference: CMS164v7) 18 years or older at the beginning of the measurement period Currently diagnosed with ischemic vascular disease (IVD), diagnosed with acute myocardial infarction (AMI), or had a coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the twelve months prior to the beginning of the measurement period Had at least one medical encounter during measurement period Exclusion: using an anticoagulant medication in the year prior to the end of the measurement period or in hospice care during the measurement period 	Patients with documentation demonstrating that Aspirin or other antiplatelet medication was being used during the year prior to the end of the measurement period						

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Asthma Medication Ratio			<ul style="list-style-type: none"> Continuously enrolled PHP members between 5 and 64 years of age at the end of the measurement period Diagnosed with persistent asthma During the past two years, had either <ol style="list-style-type: none"> at least four face-to-face medical encounters with different dates of service and at least two asthma medication dispensing events for any controller or reliever medication OR at least four asthma medication dispensing events for any controller or reliever medication Exclude patients who had no asthma medications (controller or reliever) dispensed during the year prior to the end of the measurement period Denominator exclusion if patient had any of the following diagnoses before the end of the measurement period: emphysema, other emphysema, chronic obstructive pulmonary disease, obstructive chronic bronchitis, chronic respiratory conditions due to fumes/vapors, Cystic Fibrosis, or acute respiratory failure 	<ul style="list-style-type: none"> A denominator patient qualifies for the numerator if the medication ratio is equal to or above 0.50 The medication ratio is calculated by summarizing the units of asthma controller medications and dividing it by the sum of units of total asthma medications (total = controller + reliever medications) See QIP measure specifications for a list of asthma controller medications and asthma reliever medications. The total asthma medication set is a combination of all asthma controller medications plus all asthma reliever medications. See QIP measure specifications for the definition of asthma medication dispensing events for oral, inhaler and injected medications. Each event is associated with a number of units of medication depending on the type of medication and the supply dispensed at the event 				
Early Entry Into Prenatal Care	(No eCQM reference) Patients who had any of the following within the measurement period: <ul style="list-style-type: none"> Prenatal intake Prenatal visit Birth of child 	Number of prenatal care patients who entered treatment during the first trimester For measurement purposes, the measure is reported in two ways: <ul style="list-style-type: none"> Denominator is broken down into age categories Denominator is broken down into trimester of entry into care and if the patient was first seen for prenatal care by the grantee or another provider 						

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Birth Weight From Deliveries	<ul style="list-style-type: none"> (No eCQM reference) Live births to prenatal care patients during the measurement period 	<ul style="list-style-type: none"> Infants in the denominator born with a birthweight of under 2500 grams. Note that this measure focuses on the negative outcome For measurement purposes, the denominator is broken down into birth weight categories (<1500g, 1500 to 2499g, and >=2500g) by race and ethnicity 						
Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents	<ul style="list-style-type: none"> (Reference: CMS155v9) Between 3 and 16 years at the beginning of the measurement period Had at least one medical encounter during measurement period Exclusion: pregnant or receiving hospice care during the measurement period 	<ul style="list-style-type: none"> Patients with all of the following documented in the year prior to the end of the measurement period: <ul style="list-style-type: none"> Documented height, weight and body mass index (BMI) percentile Patient (or patient's parent, as appropriate) has been counseled on nutrition Patient (or patient's parent, as appropriate) has been counseled on physical activity 						
Counseling for Nutrition for Children and Adolescents			<ul style="list-style-type: none"> Continuously enrolled PHP members between 3 and 17 years of age at the end of the measurement period Had a least one outpatient visit with a PCP or an OB/GYN during the measurement period Exclusion: pregnant during the measurement period 	<ul style="list-style-type: none"> Patients with evidence that counseling for nutrition or referral for nutrition education was documented at least once during the measurement period 				
Counseling for Physical Activity for Children and Adolescents			<ul style="list-style-type: none"> Continuously enrolled PHP members between 3 and 17 years of age at the end of the measurement period Had a least one outpatient visit with a PCP or an OB/GYN during the measurement period Exclusion: pregnant during the measurement period 	<ul style="list-style-type: none"> Patients with evidence that counseling for physical activity or referral for physical activity was documented at least once during the measurement period 				
Adult Body Mass Index (BMI) Screening and Follow-Up Plan	<ul style="list-style-type: none"> (Reference: CMS69v9) 18 years or older on the date of the last medical visit in the measurement period Had at least one medical encounter during measurement period Exclusions: pregnant during measurement period; received palliative or hospice care during measurement period; no BMI in measurement period due to patient refusal or an acceptable medical reason 	<ul style="list-style-type: none"> Patients with a BMI documented at last visit in the measurement period or within 12 months of the last visit When the most recent BMI is outside parameters, a follow up plan is documented at that visit or within 12 months of that visit. Normal weight parameters: BMI >= 18.5 and BMI < 25 kg/m2 					<ul style="list-style-type: none"> (Reference: CMS69v9) 18 years or older on the date of the last medical visit in the measurement period Had at least one medical encounter during measurement period Exclusions: pregnant during measurement period; received palliative or hospice care during measurement period; no BMI in measurement period due to patient refusal or an acceptable medical reason 	<ul style="list-style-type: none"> Patients with a BMI documented at last visit in the measurement period or within 12 months of the last visit When the most recent BMI is outside parameters, a follow up plan is documented at that visit or within 12 months of that visit. Normal weight parameters: BMI >= 18.5 and BMI < 25 kg/m2

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Tobacco Use Screening and Cessation Intervention	<ul style="list-style-type: none"> (Reference: CMS138v9) 18 years of age or older at the beginning of the measurement period Had at least two medical encounters or at least one preventive visit during the year prior to the end of the measurement period Exclusions: limited life expectancy, or no tobacco screening or intervention due to a medical reason 	<ul style="list-style-type: none"> Patients who had a tobacco assessment within one year prior to the end of the measurement period If the patient was found to be a tobacco user during the most recent assessment, the patient had at least one of the following between that date and the end of the measurement period: <ul style="list-style-type: none"> Tobacco use cessation counseling Smoking cessation agents (active medication or order) 					<ul style="list-style-type: none"> 18 years of age or older at the beginning of the measurement period Had at least two medical encounters or at least one preventive visit during the year prior to the end of the measurement period Exclusions: limited life expectancy, or no tobacco screening or intervention due to a medical reason 	<ul style="list-style-type: none"> Patients who had a tobacco assessment within one year prior to the end of the measurement period If the patient was found to be a tobacco user during the most recent assessment, the patient had at least one of the following between that date and the end of the measurement period: <ul style="list-style-type: none"> Tobacco use cessation counseling Smoking cessation agents (active medication or order)
Screening for Depression and Follow-Up Plan	<ul style="list-style-type: none"> (Reference: CMS2v10) 12 years of age or older at the beginning of the measurement period Had at least one medical encounter during measurement period Exclusions: patients with an active diagnosis for Depression or Bipolar Disorder beginning before the first encounter in the measurement period; patients who refuse to participate; patients in an urgent or emergent situation 	<ul style="list-style-type: none"> Patients who were screened for clinical depression in the past year using an age appropriate standardized tool If the screen was positive, a follow-up plan was documented on the same date as the positive screen 					<ul style="list-style-type: none"> 12 years of age or older at the beginning of the measurement period Had at least one medical encounter during measurement period Exclusions: patients with an active diagnosis for Depression or Bipolar Disorder beginning before the first encounter in the measurement period; patients who refuse to participate; patients in an urgent or emergent situation 	<ul style="list-style-type: none"> Patients who were screened for clinical depression in the past year using an age appropriate standardized tool If the screen was positive, a follow-up plan was documented on the same date as the positive screen
Depression Remission at Twelve Months	<ul style="list-style-type: none"> (Reference: CMS159v9) 12 years of age or older at the beginning of the measurement period Had at least one medical encounter during measurement period Diagnosis of major depression or dysthymia An initial PHQ-9 score greater than nine during the "index event" outpatient encounter Exclusion: patients who also have a diagnosis of bipolar disorder, personality disorder emotionally labile, schizophrenia, psychotic disorder or pervasive developmental disorder; patients who are in palliative care, hospice, a permanent nursing home, or who have died 	<ul style="list-style-type: none"> Patients who achieved remission at twelve months. This is defined as having a PHQ-9 with a score of less than five at twelve months (plus or minus 60 days) after the date of the "index event" 						

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
HIV Screening	<ul style="list-style-type: none"> • (Reference: CMS349v3) • Between 15 and 65 years of age at the beginning of the measurement period • Had at least one medical encounter during measurement period • Exclusion: Patients diagnosed with HIV prior to the beginning of the measurement period 	Patients with documentation of an HIV test performed on or after their 15th birthday and before their 66th birthday						
HIV Linkage to Care	<ul style="list-style-type: none"> • (No eCQM reference) • Patients whose first ever HIV diagnosis was made by health center staff between December 1 (of the year prior to the measurement year) and November 30 (of the current measurement year) • Had at least one medical encounter in the current measurement year or the year prior 	Patients who received treatment for HIV within 30 days of the new HIV diagnosis						
Well-Child Visits in the First 15 Months of Life			<ul style="list-style-type: none"> • Continuously enrolled PHP members who turned 15 months old during the measurement period • Exclusion: was in hospice during the measurement period 	<ul style="list-style-type: none"> • Patients with at least six well-child visits with a PCP by the date of age 15 months (which is child's first birthday plus 90 days) • There must be at least 14 days between each well-child date of service 	Patients who had at least one visit after 2 months of age and who turn 15 months old during the measurement period	Patients with at least six well-child visits with a PCP by the date of age 15 months (which is child's first birthday plus 90 days)		
Child and Adolescent Well-Care Visits			<ul style="list-style-type: none"> • Continuously enrolled PHP members between 3 and 17 years of age at the end of the measurement period • Exclusion: Patients who were in hospice care during the measurement period 	Patients with at least one well-child visit with a PCP or OB/GYN during the measurement period				

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)		
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	
Childhood Immunization Status	<ul style="list-style-type: none"> (Reference: CMS117v9) Patients who had their second birthday within measurement period Had at least one medical encounter during measurement period Exclusion: Patients who were in hospice care during the measurement period 	<ul style="list-style-type: none"> Patients with documentation of a full immunization on or before the second birthday A full immunization is all of the following: 4 diphtheria, tetanus and acellular pertussis (DTaP); 3 polio (IPV), 1 measles, mumps and rubella (MMR); 3 or 4 haemophilus influenza type B (HiB) depending on type; 3 hepatitis B (Hep B); 1 chicken pox (VZV); 4 pneumococcal conjugate (PCV); 1 hepatitis A (Hep A); 2 or 3 rotavirus (RV) depending on type; and 2 influenza (flu) vaccines Alternately, by their second birthday, a history of illness for the vaccine-preventable disease, a seropositive test result for the disease, or an allergic reaction to the vaccine can also count toward the numerator for that specific vaccine <p>Also count in the numerator the following conditions for each specific vaccine:</p> <ul style="list-style-type: none"> DTaP: encephalopathy IPV: anaphylactic reaction to streptomycin, polymyxin B, or neomycin Influenza, MMR, or VZV vaccines: cancer of lymphoreticular or histiocytic tissue, multiple myeloma, or 	<ul style="list-style-type: none"> Continuously enrolled PHP members who had their second birthday within measurement period Exclusion: Patients who were in hospice care during the measurement period 	<ul style="list-style-type: none"> Patients with documentation of a full immunization between the minimum age and the second birthday as specified in the QIP documents A full immunization is all of the following: 4 diphtheria, tetanus and acellular pertussis (DTaP); 3 polio (IPV), 1 measles, mumps and rubella (MMR); 3 or 4 haemophilus influenza type B (HiB) depending on type; 3 hepatitis B (Hep B); 1 chicken pox (VZV); 4 pneumococcal conjugate (PCV); 1 hepatitis A (Hep A); 2 or 3 rotavirus (RV) depending on type; and 2 influenza (flu) vaccines Alternately, by their second birthday, a history of illness for the vaccine-preventable disease, a seropositive test result for the disease, or an allergic reaction to the vaccine can also count toward the numerator for that specific vaccine <p>Also count in the numerator the following conditions for each specific vaccine:</p> <ul style="list-style-type: none"> DTaP: encephalopathy IPV: anaphylactic reaction to streptomycin, polymyxin B, or neomycin Influenza, MMR, or VZV 					
Immunizations for Adolescents			<ul style="list-style-type: none"> Continuously enrolled PHP members who had their 13th birthday within measurement period Exclusions: Hospice care in the measurement period or an anaphylactic reaction to any of the vaccines. Furthermore, encephalopathy due to vaccination is an exclusion for the Tdap vaccine. 	<ul style="list-style-type: none"> A full immunization is all of the following: at least one dose of meningococcal vaccine between the 11th and 13th birthdays; at least one tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap) between the 11th and 13th birthdays; and the HPV vaccination series completed between the 11th and 13th birthdays The HPV series can be two doses given at least 146 days apart or three doses with different dates of service Alternately, by the 13th birthday, a seropositive test result for the disease can also count toward the numerator for that specific vaccine 					

Measure Name	UDS/BPHC (2021)		QIP/Partnership (2021)		PIP (2021)		PHASE (2021)	
	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator	Denominator	Numerator
Dental Sealants for Children Between 6 – 9 Years	<ul style="list-style-type: none"> • (Reference: CMS277v0) • Between 6 and 9 years of age at the beginning of the measurement period • Had at least one oral assessment or comprehensive or periodic oral evaluation during a dental visit in the past year • Found to be at moderate to high risk for caries in the past year • Exclusion: patients with all first permanent molars non-sealable 	Patients who received a sealant on a permanent first molar tooth in the year prior to the end of the measurement period						